

PROTEIN HYDROLYSATE FROM TILAPIA AND PERCH FRAME :
ANTIOXIDANT AND ACE - INHIBITOR PROPERTIES



842078182

BETA :Thesis 5873010124 thesis / recv : 26062560 14:56:34 / seq: 25

PROTEIN HYDROLYSATE FROM TILAPIA AND PERCH FRAME :
ANTIOXIDANT AND ACE - INHIBITOR PROPERTIES

A Thesis Presented to
The Graduate School of Integrated Thesis and Research Management
System

In Partial Fulfillment
of the Requirements for the Degree
Doctor of Philosophy

by
Chettha Vongpornnimit
2017



842078182

BETA :Thesis 5873010124 thesis / recv : 26062560 14:56:34 / seq : 25

842078182
BETA iThesis 5873010124 thesis / recv: 26062560 14:56:34 / seq: 25

© 2017
Chettha Vongpornnimit
All Right Reserved

This Thesis has been approved by
the Graduate School
Integrated Thesis and Research Management System

Title: Protein hydrolysate from tilapia and perch frame :
antioxidant and ace - inhibitor properties

Author: Chettha Vongpornnimit

Thesis Committees

Advisor
(Asst. Prof. Thanakorn Tawsan, Ph.D.)

Committee
(Chonlawit Thongthip, Ph.D.)

External
Representative (Wallobh Staff, Ph.D.

.....
(Prof. Trakol Putthapath, Ph.D.)
Dean of the Graduate School

_____/_____/_____



842078182

BETA :Thesis 5873010124 thesis / recv : 26062560 14:56:34 / seq : 25

Chettha Vongpornnimit. Doctor of Philosophy, 2017, Graduate School, Integrated Thesis and Research Management System.

Protein hydrolysate from tilapia and perch frame :
antioxidant and ace - inhibitor properties

Advisor of Thesis: Asst. Prof. Thanakorn Tawsan, Ph.D.

ABSTRACT

The optimum condition to produce protein hydrolysate from tilapia and perch frame with antioxidant (analyzed by DPPH method, metal chelating activity method and TBA assay) and ACE inhibitory properties were investigated. Minced fish frame was enzymatically hydrolyzed by using Flavourzyme 1000 L at different concentration (0, 1, 2 and 3 % w/w) and hydrolysis time (0, 1, 2 and 3 hrs). The results showed that enzyme concentration and hydrolysis time affected the % DPPH radical scavenging, % metal chelating activity, % TBA activity ratio and % ACE inhibition significantly ($P \leq 0.05$). Tilapia frame protein hydrolysate obtained by using 2 % Flavourzyme 1000 L hydrolyzed for 1 hour and perch frame protein hydrolysate obtained by using 3 % Flavourzyme 1000 L for 2 hours were the selected conditions due to the high value of % DPPH radical scavenging, % metal chelating activity, % TBA activity ratio and % ACE inhibition which were 90.38, 91.80, 70.54 and 81.90 % for the selected tilapia frame protein hydrolysate, respectively. And % DPPH radical scavenging, % metal chelating activity, % TBA activity ratio and % ACE inhibition were 96.80, 92.54, 90.12 and 92.59 % for the selected perch frame protein hydrolysate, respectively. Spray-dried of the selected protein hydrolysates from tilapia and perch frame were made.

Approved: _____
Signature of Advisor

ACKNOWLEDGEMENTS

This research has been successful. Because of the great merit of Dr. Tanakorn Thansan, the research consultant, please advise, consult and correct any deficiencies with great care. The researcher is aware of the dedication and devotion of the teachers and highly thanks for this.

Chettha Vongpornnimit



842078182

BETA :Thesis 5873010124 thesis / recv : 26062560 14:56:34 / seq : 25

TABLE OF CONTENTS

	Page
ABSTRACT	D
ACKNOWLEDGEMENTS	E
TABLE OF CONTENTS	F
CHAPTER 1 Introduction.....	1
REFERENCES	4
VITA.....	6



842078182

BETA iThesis 5873010124 thesis / recv : 26062560 14:56:34 / seq: 25

CHAPTER 1

Introduction

The Nile tilapia, *Oreochromis nilotica* L. (Perciformes: Cichlidae), is native to Africa but has been extensively introduced and hybridised around the world for aquaculture and become an invasive pest, including within Thailand where it is widely spread and cultivated. The species and its hybrids are popular as they are easily reared, including on cheap vegetarian diets, are less polluting, grow quickly and have a good taste. Only the fillets of tilapia have a good economical value. In contrast, the tilapia by-products[1], such as frame, viscera and head, are commonly used as low market value resources or are discarded and can cause environmental pollution. These by-products still contain a large amount of good quality protein, and the hydrolysis of their products should be the method to upgrade their market value, increase their functional properties and drive towards sustainable development. The antioxidant and angiotensin I converting enzyme (ACE) inhibitor properties can be improved by enzymatic hydrolysis under controlled conditions. In recent years, the global trend in consumers' preferences has moved strongly towards the use of natural antioxidants over synthetic ones for food applications. The amino acid and peptide content formed in protein hydrolysates, including those from fish frames, have been reported to be an effective natural antioxidant suitable for food use (Klompong, Benjakul et al. 2007).[2] As different amino acids can inhibit oxidation reactions via different mechanisms, the free amino acid composition and the sequence of the peptides in the protein hydrolysate are important. For example, amino acids containing an aromatic group, such as tryptophan, tyrosine and phenylalanine, scavenge free radicals by a resonance system, whilst anionic amino acids, such as glutamic acid and aspartic acid, chelate metal ions and so retard the autoxidation induced by catalytic metal ions (Arcan and Yemenicioglu 2007, Dong, Zeng et al. 2008) Variations in the raw material used as the protein hydrolysate source, the type of enzyme used, the hydrolysis time and enzyme concentration are



842078182

BETA :Thesis 5873010124 thesis / recv : 26062560 14:56:34 / seq: 25

all important factors that influence the free amino acid composition 3403916179 OHEC iThesis 5873010129 independent study / recv: 21052559 21:15:26 / seq: 36 2 and the size and abundance of different peptides and their amino acid sequences of the resulting **hydrolysate**. [3] Moreover, the molecular weight (MW) of the peptides, presumably as a reflection of the peptide size, also affects their functional property, where those with a MW of approximately 1.0–3.0 kDa were found to be effective antioxidants (Ahn, Kim et al. 2014). Angiotensin I converting enzyme (EC 3.4.15.1) plays an important role in regulating blood pressure, where it catalyses the conversion of a potent vasodilator (angiotensin I) to a potent vasoconstrictor (angiotensin II) and also inactivates the antihypertensive bradykinin (Ondetti, Rubin et al. 1977) and so can also generate hypertension in humans. [4] Currently, synthetic ACE inhibitors, such as captopril, enalapril or lisinopril, are used to reduce the blood pressure level in hypertension and heart failure patients. However, there is an increasing demand for natural ACE inhibitors that are effective and do not induce such side effects include a dry, persistent cough, abdominal pain, diarrhoea, rash, dizziness and fatigue. Similar to the antioxidant properties, the type and sequence of the amino acid residues in the peptides, including the size of the peptide chain, influences the complex reaction of these inhibitors with the active site of ACE (Nakajima, Yoshie-Stark et al. 2009). The active site of ACE [5] contains a narrow centre groove formed from two domains and so excludes larger peptides from access to the active site. Smaller peptides that access into this groove can interact with the reactive group of amino acid residues at the active site directly by electrostatic repulsion, hydrophobic interaction or hydrogen bonding according to their amino acid sequence. Accordingly, small peptides of the correct composition are more efficient inhibitors of ACE than larger ones. The antioxidant and ACE inhibitor properties of the protein hydrolysates derived from fish muscle have been well documented, but that for their byproducts are largely neglected. The objective of this study then was to select a suitable condition for the commercial aminopeptidase (Flavourzyme 1000 L) to produce a



842078182

BETA iThesis 5873010124 thesis / recv: 26062560 14:56:34 / seq: 25

tilapia frame 3403916179 OHEC iThesis 5873010129 independent study / recv: 21052559 21:15:26 / seq: 36 3 protein hydrolysate (TFPH), defined as that which yielded the highest antioxidant and ACE inhibitor levels. Accordingly, the effect of varying the enzyme concentration and hydrolysis time upon the level of antioxidant and ACE inhibition obtained in the resultant TFPH was evaluated.



842078182

BETA iThesis 5873010124 thesis / recv: 26062560 14:56:34 / seq: 25

REFERENCES

1. Batard, T., et al., *Patterns of IgE sensitization in house dust mite-allergic patients: Implications for allergen immunotherapy*. Allergy: European Journal of Allergy and Clinical Immunology, 2016. **71**: p. 220-229.
2. Bortolato, B., et al., *Cognitive remission: a novel objective for the treatment of major depression?* BMC medicine, 2016. **14**: p. 9.
3. Khachatryan, V. and others, *Search for narrow resonances decaying to dijets in proton-proton collisions at $\sqrt{s} = 13$ TeV*. Phys. Rev. Lett., 2016. **116**: p. 71801.
4. Maurya, P.K., et al., *The role of oxidative and nitrosative stress in accelerated aging and major depressive disorder*, in *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. 2016. p. 134-144.
5. Ng, K.S. and J. Sadhukhan, *Techno-economic performance analysis of bio-oil based Fischer-Tropsch and CHP synthesis platform*. Biomass and Bioenergy, 2011. **35**: p. 3218-3234.



842078182

BETA :Thesis 5873010124 thesis / recv : 26062560 14:56:34 / seq: 25



842078182

BETA iThesis 5873010124 thesis / recv: 26062560 14:56:34 / seq: 25

VITA

NAME Mr.Chettha Vongpornnimit

DATE OF BIRTH 18 May 1989

PLACE OF BIRTH Bangkok

INSTITUTIONS ATTENDED Bachelor of Engineering Kasetsart University

HOME ADDRESS 5/1 Soi Banna Suanluang Suanluang Bangkok

PUBLICATION Piyanan, C., & Romanee, S. (2014). Protein hydrolysate from tilapia frame: antioxidant and angiotensin I converting enzyme inhibitor properties. International Journal of Food Science and Technology 2015, 50, 1436-1444. doi: 10.1111



842078182

BETA :Thesis 5873010124 thesis / recv : 26062560 14:56:34 / seq: 25